

TRIANUM-P

BIOLOGICAL FUNGICIDE



ACTIVE INGREDIENT

Trichoderma harzianum Rifai strain KRL-AG2*

3.65%

OTHER INGREDIENTS

96.35%

TOTAL

100.00%

*Contains at least 1.0×10^7 colony forming units per gram dry weight.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container under refrigerated conditions. Short periods at room temperatures above 75°F will not affect performance. Do not store near food or feed commodities. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL

Do not contaminate food or feed by disposal. Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Completely empty bag into the application equipment by shaking and tapping sides and bottom to loosen clinging particles, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration.

NET WEIGHT

1.1 lbs (500 gram)

EPA Reg. No. 89635-3

EPA Est. No. 63119-NLD-001

PRODUCT NUMBER

12903

MANUFACTURING DATE/
EXPIRATION DATE /
LOT NUMBER

see packaging

MANUFACTURED BY

Koppert Biological Systems, Inc.
1502 Old US 23
Howell, Michigan 48843
U.S.A.



8 719747 200216

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixer/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization. Follow manufacturer's directions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE from other laundry.

USER SAFETY RECOMMENDATIONS

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. For any requirement specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted - entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is protective eyewear, coveralls, waterproof gloves, shoes, and socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants or farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled.

GENERAL INFORMATION

Trianum-P Biological Fungicide is a preventative biological fungicide for control of plant diseases. The active ingredient is a microbe, *Trichoderma harzianum* Strain T-22, which when applied to seeds, transplants or other propagative material, or to soil or planting mixes, grows onto plant roots as they develop and provides protection against plant root pathogens including *Pythium*, *Rhizoctonia*, *Fusarium*, *Cylindrocladium* and *Thielaviopsis*. Trianum-P Biological Fungicide can be used alone or in conjunction with certain chemical fungicides; consult Trianum-P Biological Fungicide compatibility chart, tank mix compatibility charts below or the company for more information.

This product should not be tank mixed with chemicals that contain the following active ingredients: benzoyl, imazalil, propiconazole, tebuconazole, and triflumizole. Do not apply Trianum-P Biological Fungicide immediately before these pesticides are used. See specific instructions for tank mixing. Where early season seed rot and seedling diseases are expected, use chemically treated seed or other appropriate measures for stand establishment and Trianum-P Biological Fungicide for root disease control.

Note: Trianum-P Biological Fungicide contains live spores of a microbe that must be used prior to disease onset. Trianum-P Biological Fungicide becomes active in soil or on plants when temperatures are above 50°F and is not effective while temperatures remain cold. Trianum-P Biological Fungicide can be applied to sterilized or fumigated soil but must be applied after sterilization or fumigation.

This biological fungicide is for use in soil applications (drench, in soil furrow, potting soil and broadcast), and seed treatments in or on all raw agricultural commodities, food and fiber crops.

Trianum-P Biological Fungicide is for use in soil applications (drench, in soil furrow, and potting soil), and seed treatments on ornamentals, landscape plants, turf, and ornamental trees, including tree seedlings for transplanting into the forest.

If Trianum-P Biological Fungicide is mixed with water to make a slurry, dip or suspension, use immediately. If not, keep refrigerated after mixing with water.

NOTE: DO NOT APPLY to sugarcane, peach, rice, mushrooms, kiwi, tobacco, barley, oats, wheat, lemon, apple, and chickpea. Not for use on aquatic crops.

For food commodities: Use in chemigation and irrigation systems is limited to greenhouse flood, drip, furrow, micro-irrigation, and ebb and flow applications with NO OVERHEAD SPRAY. See footnotes for specific directions concerning each use pattern.

APPLY VIA GROUND APPLICATION ONLY

CROPS	USE	RATE
Agronomic Row or Other Field Crops: Buckwheat, Beans (soybean, snap, dry), Corn (grain, seed, sweet corn, silage, popcorn, high oil), Cotton, Canola, Peas (dry, succulent), Safflower, Sunflower	Planter Box (on-site) Commercial seed treatment	1.0 - 10.0 oz. / cwt. seed 0.00035 - 26.43 lbs. / cwt. seed
Alfalfa Hay and Forage Crops: Alfalfa, Clover, Vetch, Trefoil	Planter Box (on-site) Commercial seed treatment	1.0- 10;0 oz. / cwt. seed 0.00035 - 26.43 lbs. / cwt. seed
Berries and Small Fruits: Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries, Strawberries, Grapes	Cuttings/bare root Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution *Greenhouse chemigation	0.25- 5.0 lbs. / 5 gal or dip into dry powder. 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 32.0 oz. / 100 gal
Bulb Crops: Garlic, Leeks, Onions, Shallots, Ornamental Bulbs	Dust (pre-plant)	0.03 - 3.0 lbs. / cwt. seed
Citrus Fruits: Citrus Hybrids, Grapefruit, Kumquat, Limes, Oranges, Pummelos	Cuttings or bare-roots Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution *Greenhouse chemigation	0.25 - 5.0 lbs. / 5 gal or dip into dry powder. 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 32.0 oz. / 100 gal
Cucurbit Vegetables: Cucumbers, Melons, Gourds Pumpkins, Squash	Planter Box (on-site) Commercial seed treatment Greenhouse soil drench In-furrow spray or transplant starter solution *Greenhouse chemigation	1.0 -10.0 oz. / cwt. seed 0.00035 - 26.43 lbs./ cwt. seed 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 32.0 oz. / 100 gal
Flowers, Bedding Plants, and Ornamentals	Cuttings or bare-roots Commercial seed treatment Greenhouse soil drench Nursery soil drench *Greenhouse chemigation	0.25 - 5.0 lbs. / 5 gal or dip into dry powder. 0.088 - 17.62 lbs. / cwt. seed 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal

CROPS	USE	RATE
Fruiting Vegetables: Eggplant, Sweet and Hot Peppers, Tomatillos, Tomatoes	Commercial seed treatment Greenhouse soil drench In-furrow spray or transplant starter solution *Greenhouse chemigation	0.00035 - 26.43 lbs. / cwt. seed 1.0 c- 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 32.0 oz. / 100 gal
Herbs, Spices, and Mints	Commercial seed treatment Greenhouse soil drench starter solution *Greenhouse chemigation In-furrow spray or transplant	0.00035 - 26.43 lbs. / cwt. seed 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal
Hydroponic Crops: Cucumbers, Tomatoes, Lettuce, Herbs and Spices	Greenhouse soil drench *Greenhouse chemigation	1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal
Leafy Vegetables and Cole Crops: Arugula, Celery, Chervil, Endive, Fennel, Lettuce (head and leaf), Parsley, Radicchio, Rhubarb, Spinach, Swiss Chard, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Asparagus	Cuttings or bare-roots Commercial seed treatment In-furrow spray or transplant starter solution Greenhouse soil drench *Greenhouse Chemigation	0.25 - 5.0 lbs. / 5 gal or dip into dry powder. 0.00035 - 26.43 lbs. / cwt. seed 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 32.0 oz. / 100 gal
Legume Vegetable and Fields Crops: Snap and Dry Beans, Lentils, Succulent and Dry Peas, Peanuts, Soybeans	Planter Box (Onsite) Commercial seed treatment	1.0 - 10.0 oz. / cwt. seed 1.0 - 3.0 tbsp. / gal
Pome Fruit: Pears, Quince	Greenhouse soil drench Nursery soil drench starter solution *Greenhouse chemigation In-furrow spray or transplant	1.0 - 32.0 oz. / 100 gal 1-0- 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal
Root Crops: Beets, Sugar beets, Red Beets, Carrots, Celeriac, Chicory, Horseradish, Parsnip, Radish, Rutabaga, Salsify, Turnips	Planter Box (Onsite) Commercial seed treatment	1.0 - 10.0 oz. / cwt. seed 0.00035 - 26.43 lbs. / cwt. seed

CROPS	USE	RATE
Shade house and Outdoor Nursery Crops: Deciduous Trees (Maples, Oak, etc.), Ornamentals, Grapes, Citrus, Pine	Cuttings or bare-roots Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution *Greenhouse chemigation	0.25 - 5.0 lbs. / 5 gal or dip into dry powder. 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 32.0 oz. / 100 gal
Small Grains: Rye, Wheat, Sorghum, Millet	Planter Box (Onsite) Commercial seed treatment	1.0 - 10.0 oz. / cwt. seed 0.00035 - 26.43 lbs. / cwt. seed
Stone Fruit: Apricots, Cherries, Nectarines, Peaches, Plums, Prunes	Cuttings or bare-roots Greenhouse soil drench Nursery soil drench Cuttings or bare-roots *Greenhouse chemigation In-furrow spray or transplant	0.25 - 5.0 lbs. 15 gal or dip into dry powder. 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 3.0 tbsp. / gal 1.0 - 32.0 oz. / 100 gal
Tree Nuts: Almonds, Beech Nuts, Brazil Nuts, Butternuts Cashews, Chestnuts, Filberts, Hickory Nuts, Macadamia Nuts, Pecans, Pistachios, Walnuts	Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution *Greenhouse chemigation	1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / 100 gal 1.0 - 32.0 oz. / acre 1.0 - 32.0 oz./ 100 gal
Tuber Crops: Potatoes, Sweet Potatoes, Yams, Jerusalem Artichoke, Cassava, Ginger	Planter Box (Onsite) In-furrow spray or transplant starter solution	0.03 -3.0 oz. / cwt. seed 1.0 - 32.0 oz. / acre

* Application via greenhouse chemigation is limited to flood, dip, furrow, micro-irrigation, and ebb and flow systems. Do not apply product when above-ground harvestable food commodities are present. Refer to Chemigation section for specific directions.

SEED TREATMENT FOR TRUE SEED CROPS

ONSITE APPLICATION TO SEED: Trianum-P Biological Fungicide is applied to seeds at the rate of 1.0 - 10.0 ounces per hundredweight (oz. / cwt.) for protection against root diseases. For example, for large, smooth seeds such as soybean or dry bean, apply 1.0 - 10.0 oz./cwt. of seed. For smaller or rougher seed such as peas, or corn, apply 1.0 - 8.0 oz. / cwt. For sweet corn, apply 1.0 - 10.0 oz/cwt. Trianum-P Biological Fungicide can be applied in sufficient water to coat seeds. For maximum seed protection, especially in cold soils, apply Trianum-P Biological Fungicide to commercially treated seed such as seed treated with Captain, Apron and or Demosan for stand establishment.

Do not use treated seed for food or feed purposes or process for oil.
Treat only those seeds needed for immediate use, minimizing the interval between treatments and planting. Do not store excess treated seeds beyond planting time.

COMMERCIAL SEED TREATMENT

Apply Trianum-P Biological Fungicide as slurry, a coating, in a pellet, or during seed priming. See table below.

NOTE: This product does not contain a dye and is not covered by an appropriate tolerance exemption, or other clearance under the Federal Food, Drug and Cosmetic Act. To comply with 40 CFR §153.155, all seeds treated commercially with this product, must be colored with an EPA-approved colorant of suitable color to prevent accidental use as food for man or feed for animals.

The Federal Seed Act requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with Trichoderma harzianum Rifai strain KRL-AG2. Do not use for food, feed or oil purposes."

AGRICULTURAL CROPS

SEED SIZE (#SEEDS/OZ.)	GRAMS OF TRIANUM-P PER LB. OF SEED	OUNCES OF TRIANUM-P PER LB. OF SEED
Large (1-100) e.g. peanuts, green & dry beans, field corn	0.0016 to 0.32 g	.0016 to 0.32 oz.
Medium (100-1000) e.g. sweet corn, soybeans, sorghum	0.08 to 2.4 g	.0028 to 0.0847 oz.
Small (1000-10,000) e.g. cabbage, cucumbers, sugar beets	0.4 to 16 g	0.141 to 0.564 oz.
Fine (10,000-100,000) e.g. tomatoes	1.6 to 120 g	0.9564 to 4.23 oz.

ORNAMENTAL CROPS

SEED SIZE (#SEEDS/OZ.)	GRAMS OF TRIANUM-P PER LB. OF SEED	OUNCES OF TRIANUM-P PER LB. OF SEED
Small (1,000 - 10,000) e.g. Echinacea, Cosmos	0.4-16 g	0.0141 to 0.564 oz.
Fine (10,000 - 100,000) e.g. Texas bluegrass, ryegrass	1.6 - 120 g	0.0564 to 4.23 oz.
Very fine (100,000 - 500,000) e.g. bentgrass	8.0 - 120 g	0.282 to 2.82 oz.

SEED TREATMENT FOR VEGETATIVELY PROPAGATED CROPS, INCLUDING POTATOES, OTHER ROOT, TUBER AND BULB VEGETABLES

For planting or storage, treat at 0.03 - 3.0 ounces Trianum-P Biological Fungicide to 100 lbs (1 cwt) of bulbs or cut potato seed pieces. Dip bulbs, tubers or cut potato seed pieces in a suspension of 1.0 - 3.0 lbs. of Trianum-P Biological Fungicide in 20 gallons of water.

For potatoes, consult your Koppert Biological Systems Representative for more information. All surfaces, knives, and other equipment used to cut and plant potatoes should be thoroughly sterilized before cutting and planting and at regular intervals. The cut and treated seed pieces may be held for a week or more at cool temperatures, 45 - 50°F, and high relative humidity to promote suberization or they may be planted immediately.

DIPS FOR CUTTINGS AND BARE ROOTED TRANSPLANTS

Dip cuttings, bulbs or transplants in a suspension of 1.0 - 3.0 lbs. of Trianum-P Biological Fungicide in 20 gallons of water. Plant treated cuttings, bulbs or transplants in potting mix or soil in the usual manner.

SOIL DRENCH

GREENHOUSE SOIL DRENCH

Suspend 1.0 - 32.0 ounces Trianum-P Biological Fungicide in 100 gallons of water with agitation and apply as a soil drench to greenhouse planting mixes. For seeding flats or shallow (up to 4-inch depth) beds or pots, apply at a rate of 50 - 100 gallons per 800 square feet. For deeper beds or pots, apply at a rate of 100 gallons per 400 square feet, ½ cup (4 fl. ounces) for pots with a 3-inch diameter, or 1 cup (8 fl. ounces) per 6-inch diameter pot.

Apply Trianum-P Biological Fungicide through low pressure watering nozzles such as fan nozzles or other drench watering systems applied directly to the soil. Constant agitation is required to maintain Trianum-P Biological Fungicide in suspension. Trianum-P Biological Fungicide can be tank mixed and is compatible with many commonly used fungicides,

liquid fertilizers; herbicides, insecticides and biological control products registered for use on greenhouse/ornamental plants. If tank mixes are desired, observe the most restrictive of labeling limitations and precautions of all products used in mixtures. Consult the tank mix compatibility chart below or the company for more information.

NURSERY SOIL DRENCH

Suspend 1.0 - 32.0 ounces Trianum-P Biological Fungicide in 100 gallons of water with agitation and apply as a soil drench to container nursery crops. For shallow (up to 4-inch depth) beds or pots, apply at a rate of 50 - 100 gallons per 800 square feet. For deeper beds or pots, apply at a rate of 100 gallons per 400 square feet, ½ cup (4 fl. ounces) for pots with a 3-inch diameter, or 1 cup (8 fl. ounces) per 6-inch diameter pot.

Apply Trianum-P Biological Fungicide directly to the soil through low pressure watering nozzles such as fan nozzles, or other drench watering systems, handheld sprayers or backpack sprayers. Constant agitation is required to maintain Trianum-P Biological Fungicide in suspension. Trianum-P Biological Fungicide can be tank mixed and is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products registered for use on nursery plants. If tank mixes are desired, observe the most restrictive labeling limitations and precautions of all products used in mixtures. Consult the tank mix compatibility chart below or the company for more information.

IN-FURROW SPRAY OR TRANSPLANT STARTER SOLUTION

Apply as an in-furrow spray or transplant starter solution at a rate of 1.0 - 32.0 ounces/acre in sufficient water to achieve uniform application. Maintain constant agitation. Trianum-P Biological Fungicide can be tank mixed with certain fertilizers and pesticides; consult tank mix compatibility chart below for detailed information.

TANK MIXING

Trianum-P Biological Fungicide can be tank mixed and is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products registered for use on greenhouse/ornamental plants. If tank mixes are desired, observe the most restrictive of labeling limitations and precautions of all products used in mixtures. Consult the tank mix compatibility chart below or the company for more information.

This product should not be tank mixed with chemicals that contain the following active ingredients: imazilil, propiconazole, tebuconazole, and triflumizole. Do not apply Trianum-P Biological Fungicide immediately before these pesticides are used.

This product can be mixed with the specific products, their percentages and rates for use in nursery drench, in-furrow spray or transplant starter solution, as listed in the table below in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates.

This product can not be mixed with any product containing a label prohibition against such mixing.

COMPATIBILITY FOR DRENCH, IN-FURROW SPRAY OR TRANSPLANT STARTER SOLUTION TANK MIX

NOTE: While the information presented in this table is believed to be up to date, the user must always read the label of the other products used in the tank mix to confirm application rates and dilutions.

CHEMICAL NAME	% A.I. FORMULATION	PRODUCT NAME	RATE PRODUCT	DILUTION
Captan	85% Wettable Powder	Captan 85WP	1.88 lb./Acre	0.1 oz. / gal.
Chlorothalonil	82.5% Water Dispersible Granules	Daconil Ultrex	3.7 ounces 1000 ft ²	0.56 oz. / gal.
Iprodione	23.3% Flowable	Chipco 26019 Flo	4 ounces per 1000 ft ²	0.6 oz. / gal.
Thiophanate methyl	50% Wettable Powder	Cleary's 3336 in water soluble bags	8 oz. per 1000 ft ²	1.2 oz. / gal.
Iprodione	Granules 50%, Soluble	Rovral	0.75 lb/Acre	0.04 oz. / gal.
Metalaxyl	21.3% Liquid	Subdue Maxx	0.25 ounces per 800 ft ²	0.05 oz. / gal.
Chlorpyrifos	50% Emulsifiable Liquid	Lorsban 4E	3.2 ounces per gallon	3.2 oz. / gal.

GREENHOUSE CHEMIGATION

Suspend 1.0 - 32.0 ounces Triam-P Biological Fungicide in 100 gallons of water with agitation and apply through the following systems: 1) pressurized drench (flood) or drip (trickle) systems, 2) furrow, 3) micro-irrigation such as spaghetti-tube or individual tube irrigation, 4) hand-held calibrated irrigation equipment such as the hand-held wand with injector, and 5) ebb and flow systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers or other experts.

Do not connect an irrigation system, (including greenhouse system), used for pesticide application to a public water system unless the pesticide safety systems for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

GREENHOUSE CHEMIGATION

Requirements for Chemigation Systems Connected to Public Water Systems:

1. Public water systems means a system for the provision to the public of piped water for human consumption if such a system has at least 15 service connections or regular serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow riser of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Apply Trianium-P Biological Fungicide during the last half of the water application period. Mix Trianium-P Biological Fungicide in enough water to be able to draw through the system for the last half of the water application.
9. Apply enough water to move Trianium-P Biological Fungicide into the root zone. Amounts will vary depending on soil type and existing moisture level. Avoid applying water volumes that would cause runoff or excessive leaching.

Drip (Trickle) Chemigation and Micro-irrigation Requirements

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional inter-locking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Apply Trianium-P Biological Fungicide during the last half of the water application period. Mix Trianium-P Biological Fungicide in enough water to be able to draw through the system for the last half of the water application.

8. Apply enough water to move Trianum-P Biological Fungicide into the root zone. Amounts will vary depending on soil type and existing moisture level. Avoid applying water volumes that would cause runoff or excessive leaching.

PLANT SAFETY

Trianum-P Biological Fungicide has been tested on numerous plant varieties with no phytotoxic effects. However, since Trianum-P Biological Fungicide has not been tested on all plant varieties or in combination with all available tank mixes the manufacturer recommends testing Trianum-P Biological Fungicide on a small number of plants to check for adverse plant effects before applying to a larger number of plants.

NOTICE TO BUYER AND SELLER: Seller warrants that this product conforms to the description on the label and is reasonably fit for the purposes stated on the label when used and stored in accordance with directions under normal conditions of use. To the extent permitted by state law, this warranty does not extend to use of this product contrary to label directions or under conditions not reasonably foreseeable by the Seller, and Buyer and User assume the risk of any such use. To the extent permitted by state law, Seller disclaims all other warranties express or implied, including any warranty of fitness or merchantability. To the extent permitted by state law, Seller shall not be liable for consequential, special or indirect damages resulting from use or handling of this product and Seller's sole liability and Buyer's and user's exclusive remedy shall be limited to refund of the purchase price. This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified.

EPA Reg. No. 89635-3
EPA Est. No. 63119-NLD-001

MANUFACTURED BY

Koppert Biological Systems, Inc.
1502 Old US 23
Howell, Michigan 48843
U.S.A.